

Words Matter - Terms to Use and Avoid When Talking About Addiction

Image



This page offers background information and tips for providers to keep in mind while using person-first language, as well as terms to avoid to reduce stigma and negative bias when discussing addiction. Although some language that may be considered stigmatizing is commonly used within social communities of people who struggle with substance use disorder (SUD), clinicians can show leadership in how language can destigmatize the disease of addiction. For similar information for patients, visit NIDA's [Words Matter: Preferred Language for Talking About Addiction](#).

To learn more about the unique impact of stigma on pregnant women and mothers, visit NIDAMED's [Your Words Matter – Language Showing Compassion and Care for Women, Infants, Families, and Communities Impacted by Substance Use Disorder](#).

How to earn CME/CE credit:

- Step 1: Read all of the content below.
- Step 2: Go to the [Words Matter CME/CE Activity](#) page and complete the instructions provided under [Method of Participation and Request for Credit](#) to obtain your CE certificate.

Stigma and Addiction

What is stigma?

Stigma is a discrimination against an identifiable group of people, a place, or a nation. Stigma about people with SUD might include inaccurate or unfounded thoughts like they are dangerous, incapable of managing treatment, or at fault for their condition.

Where does stigma come from?

For people with an SUD, stigma may stem from antiquated and inaccurate beliefs that addiction is a moral failing, instead of what we know it to be—a chronic, treatable disease from which patients can recover and continue to lead healthy lives.

How does stigma affect people with SUD?

- Feeling stigmatized can reduce the willingness of individuals with SUD to seek treatment.^{1,2}
- Stigmatizing views of people with SUD are common; this stereotyping can lead others to feel pity, fear, anger, and a desire for social distance from people with an SUD.²
- Stigmatizing language can negatively influence health care provider perceptions of people with SUD, which can impact the care they provide.³

How can we change stigmatizing behavior?

- When talking to people with SUD, their loved ones, and your colleagues, use non-stigmatizing language that reflects an accurate, science-based understanding of SUD and is consistent with your professional role.
- Because clinicians are typically the first points of contact for a person with an SUD, health professionals should “take all steps necessary to reduce the potential for stigma and negative bias.”³ Take the first step by learning the terms to avoid and use.
- Use person-first language and let individuals choose how they are described.⁴ Person-first language maintains the integrity of individuals as whole human beings—by removing language that equates people to their condition or has negative connotations.⁵ For example, “person with a substance use disorder” has a neutral tone and distinguishes the person from his or her diagnosis.⁶

What else should I keep in mind?

It is recommended that “substance use” be used to describe all substances, including alcohol and other drugs, and that clinicians refer to severity specifiers (e.g., mild, moderate, severe) to indicate the severity of the SUD. This language also supports documentation of accurate clinical assessment and development of effective treatment plans.⁷ When talking about treatment plans with people with SUD and their loved ones, be sure to use evidence-based language instead of referring to treatment as an intervention.

Terms to avoid, terms to use, and why

Consider using these recommended terms to reduce stigma and negative bias when talking about addiction.

Instead of...	Use...	Because...
<ul style="list-style-type: none"> Addict 	<ul style="list-style-type: none"> Person with substance use disorder¹ 	<ul style="list-style-type: none"> Person-first language. The change shows that “has” a problem, rather than “is” a problem.⁷ The terms avoid eliciting negative associations, punitive attitudes, and individual blame.⁷
<ul style="list-style-type: none"> User 	<ul style="list-style-type: none"> Person with OUD or person with opioid addiction (when substance in use is opioids) 	
<ul style="list-style-type: none"> Substance or drug abuser 	<ul style="list-style-type: none"> Patient 	
<ul style="list-style-type: none"> Junkie 	<ul style="list-style-type: none"> Person in active use; use the person’s name, and then say “is in active use.” 	
<ul style="list-style-type: none"> Alcoholic 	<ul style="list-style-type: none"> Person with alcohol use disorder 	
<ul style="list-style-type: none"> Drunk 	<ul style="list-style-type: none"> Person who misuses alcohol/engages in unhealthy/hazardous alcohol use 	
<ul style="list-style-type: none"> Former addict 	<ul style="list-style-type: none"> Person in recovery or long-term recovery 	
<ul style="list-style-type: none"> Reformed addict 	<ul style="list-style-type: none"> Person who previously used drugs 	
<ul style="list-style-type: none"> Habit 	<ul style="list-style-type: none"> Substance use disorder 	

Instead of...	Use...	Because...
	<ul style="list-style-type: none"> ▪ Drug addiction 	<ul style="list-style-type: none"> ▪ “Habit” may undermine the seriousness of the disease.
<ul style="list-style-type: none"> ▪ Abuse 	<p>For illicit drugs:</p> <ul style="list-style-type: none"> ▪ Use <p>For prescription medications:</p> <ul style="list-style-type: none"> ▪ Misuse ▪ Used other than prescribed 	<ul style="list-style-type: none"> ▪ The term “abuse” was found to have a negative association with negative judgments and punishment.⁹ ▪ Legitimate use of prescription medication is not abuse. Abuse is defined as use other than as prescribed by the person to whom they are prescribed. Consumption outside of these parameters is misuse.
<ul style="list-style-type: none"> ▪ Opioid substitution replacement therapy ▪ Medication-assisted treatment (MAT) 	<ul style="list-style-type: none"> ▪ Opioid agonist therapy ▪ Pharmacotherapy ▪ Addiction medication ▪ Medication for a substance use disorder ▪ Medication for opioid use disorder (MOUD) 	<ul style="list-style-type: none"> ▪ It is a misconception that medications merely “substitute” one drug or “one addiction” for another. ▪ The term MAT implies that medication serves a supplemental or temporary role in treatment. “MOUD” aligns with the way other psychotropic medications are understood (e.g., antidepressants, antipsychotics), as critical tools that are part of a patient’s treatment plan.
<ul style="list-style-type: none"> ▪ Clean 	<p>For toxicology screen results:</p> <ul style="list-style-type: none"> ▪ Testing negative <p>For non-toxicology purposes:</p> <ul style="list-style-type: none"> ▪ Being in remission or recovery ▪ Abstinent from drugs ▪ Not drinking or taking drugs ▪ Not currently or actively using drugs 	<ul style="list-style-type: none"> ▪ Use clinically accurate, non-stigmatizing language the same way it would be used for other medical conditions.¹⁰ ▪ Set an example with your own language when treating patients who might use stigmatizing language. ▪ Use of such terms may evoke negative implicit cognitions.⁷
<ul style="list-style-type: none"> ▪ Dirty 	<p>For toxicology screen results:</p> <ul style="list-style-type: none"> ▪ Testing positive 	<ul style="list-style-type: none"> ▪ Use clinically accurate, non-stigmatizing language the same way it would be used for other medical conditions.⁹ ▪ May decrease patients’ sense of hope and self-efficacy for change.⁷

Instead of...	Use...	Because...
	<p>For non-toxicology purposes:</p> <ul style="list-style-type: none"> ▪ Person who uses drugs 	<ul style="list-style-type: none"> ▪
<ul style="list-style-type: none"> ▪ Addicted baby 	<ul style="list-style-type: none"> ▪ Baby born to mother who used drugs while pregnant ▪ Baby with signs of withdrawal from prenatal drug exposure ▪ Baby with neonatal opioid withdrawal/neonatal abstinence syndrome ▪ Newborn exposed to substances 	<ul style="list-style-type: none"> ▪ Babies cannot be born with addiction because addiction is a behavioral disorder—they are born manifesting a withdrawal syndrome ▪ Use clinically accurate, non-stigmatizing language the same way it would be used for other conditions.¹⁰ ▪ Using person-first language can reduce

References

1. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5937046>
2. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5854406>
3. <https://www.tandfonline.com/doi/abs/10.1080/10826084.2019.1581221?journalCode=isum20>
4. <https://www.ncbi.nlm.nih.gov/pubmed/31140667>
5. <https://apastyle.apa.org/6th-edition-resources/nonhandicapping-language>
6. <https://obamawhitehouse.archives.gov/sites/whitehouse.gov/files/images/Memo%20-%20Changing%20Federal%20Terminology%20Regrading%20Substance%20Use%20and%20Substance%20Use%20Disorders.pdf>
7. <https://www.thenationalcouncil.org/wp-content/uploads/2016/10/Substance-Use-Terminology.pdf>
8. <https://psycnet.apa.org/record/2018-44736-001>
9. <https://www.sciencedirect.com/science/article/abs/pii/S0955395909001546?via%3Dihub>
10. <https://jamanetwork.com/journals/jama/article-abstract/1838170>

November 29, 2021