

The Webinar Is Now Live

**This webinar is being
recorded**

Your audio will remain muted



Technical Information

This webinar is being recorded and archived, and will be available to all webinar participants. Please contact the webinar facilitator if you have any concerns or questions.

This training was developed under the Substance Abuse and Mental Health Services Administration's Prevention Technology Transfer Center task order. Reference # 1H79SP081018.

For training use only.

Funding for this presentation was made possible by SAMHSA grant no. 1H79SP081018. The views expressed by speakers and moderators do not necessarily reflect the official policies of HHS; nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.



Housekeeping

The **Chat and Questions** chat box to the right of the presentation is for general discussion. Use this space to...

- Interact with other participants
- Submit questions for the presenter(s)
- Respond to discussion questions

These messages are public and can be seen by everyone in the room.



Chat and Questions (Everyone)

[Empty chat area]

[Input field] [Send icon]

Everyone



Today's Audio

Please connect audio via phone OR computer - not both. 888-858-6021; Conf code: 691 584 5839

If you dialed in, you need to turn your computer speakers off or you will hear an echo.

Additionally, if you are connecting via phone, please MUTE your device to minimize

Tech Support

[Empty text area]

[Input field] [Send icon]

Housekeeping

If you have any technical difficulties, you can type your issue in the **Tech Support** chat box found below the slides.

These messages will only be seen by Tech Support staff.



Today's Audio [menu icon]

Please connect audio via phone OR computer - not both. 888-858-6021; Conf code: 691 584 5839

If you dialed in, you need to turn your computer speakers off or you will hear an echo.

Additionally, if you are connecting via phone, please MUTE your device to minimize

Tech Support [menu icon]

[Input field]

[Send icon]


Chat and Questions (Everyone) [menu icon]

[Input field]

[Send icon]

Everyone

Housekeeping



If you need to adjust your speaker audio, click the dropdown arrow next to the speaker icon and click **Adjust Speaker Volume**.



If you need to troubleshoot your computer audio you can access the **Audio Setup Wizard** by clicking the *Meeting* button.

Today's Audio information is available below the slides. If you are calling in on your phone, please leave your line muted.



Today's Audio

Please connect audio via phone OR computer - not both. 888-858-6021; Conf code: 691 584 5839

If you dialed in, you need to turn your computer speakers off or you will hear an echo.

Additionally, if you are connecting via phone, please MUTE your device to minimize

Tech Support

[Empty chat area]

Chat and Questions (Everyone)

[Empty chat area]

Everyone

Housekeeping

**Webinar
Recording**
available online
in 2-3 weeks at
bit.ly/ce-pttc

After the webinar:

Complete the
brief, 10-question
Evaluation Form



to gain access to

Resource Page

1. List of resources shared today
2. PowerPoint slides
3. Link to request Certificate of Attendance



Welcome



Central East PTTC Webinar:

Opioid and Alcohol Misuse: The Risk of Suicide

The Central East PTTC is housed at the Danya Institute in Silver Spring, MD

Renata Henry
Executive Director

Deborah Nixon Hughes
Project Director



Central East (HHS Region 3)

PTTC

Each TTC Network includes 13 centers.*



Network Coordinating Office

National American Indian and Alaska Native Center

National Hispanic and Latino Center


10 Regional Centers (aligned with HHS regions)

**The ATTC Network also includes 4 international HIV centers funded by the President's Emergency Plan for AIDS Relief.*



Central East Region

HHS REGION 3



What We Do

- **Accelerate** the adoption and implementation of evidence-based and promising treatment and recovery-oriented practices and services
- **Strengthen** the behavioral health awareness, knowledge, and skills of the health care and prevention workforce that address the needs of people with behavioral health disorders
- **Foster regional** and national alliances among culturally diverse practitioners, researchers, policy makers, funders, and the recovery community
- **Ensure** the availability and delivery of publicly available, free of charge, behavioral health training and technical assistance to health care and prevention professionals



PTTC Mission



To Strengthen **the Capacity of the Workforce to Deliver Evidence-Based Prevention Strategies and Facilitate Opportunities for Preventionists to Pursue New Collaboration Opportunities, which include Developing Prevention Partnerships and Alliances**



Central East PTTC Specialty Area

Engaging and Collaborating with Primary Care Providers for Substance Use Prevention



Eligibility

Behavioral health and health care providers, consumers, families, state and local public health systems and other stakeholders

**Consistent with
Regional, State
and Local
Needs**

No cost

Data Driven

**EBPs provided
by Subject
Matter Experts**



PTTC Focus Areas

- Opioid/harm reduction strategies
- Suicide prevention and substance use overdoses
- Youth vaping/tobacco
- Youth alcohol and suicide
- Health equity and the elimination of disparities
- Trauma
- Workforce leadership development series-
strategic planning/SPF; data-informed decision-
making; environmental strategies



Upcoming Webinars

- Social Determinants of Health: A Prevention Perspective Part 2- **May 20**
- Motivational Interviewing: African American Males- **May 27**
- Shifting the Attitudes of Law Enforcement in Addressing the Opioid Epidemic- **June 3**
- Preventing Youth Vaping Parts 1 & 2- **June 25, August 6**
- Building Capacity for a Public Health Approach to Prevention- **July 16**
- Identifying Emerging Drug Trends- **September 24**

ALL WEBINARS AT 1:00 PM ET



Other Resources in Region 3



Central East (HHS Region 3)

ATTC

Addiction Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration



Central East (HHS Region 3)

MHTTC

Mental Health Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration



Central East (HHS Region 3)

PTTC



Central East (HHS Region 3)

PTTC

Prevention Technology Transfer Center Network

Funded by Substance Abuse and Mental Health Services Administration

Opioid and Alcohol Misuse: The Risk of Suicide

May 14, 2020

Josh Esrick

PTTC Prevention Specialist

Senior Policy Analyst

Carnevale Associates, LLC

Emily Patton

Research Associate

Carnevale Associates, LLC



Presenters



Josh Esrick
Presenter



Emily Patton
Presenter



Key Objectives

- Examine the scope of suicide risk and the populations most at-risk for suicide
- Summarize the scope of substance use-involved fatalities and the impact of suicide
- Discuss the research on the connections between substance use and risk of suicide
- Overview evidence-based programs for preventing substance use and suicide, as well as other opportunities for collaboration



Central East PTTC Product

- Upcoming public product, a literature review covering:
 - Background data on the scope of opioid misuse, alcohol misuse, and risk and rates of suicide
 - How opioids and alcohol increase the risk of suicide
 - Evidence-based prevention strategies that address opioid or alcohol misuse and risk of suicide



Background on Suicide

- In 2018, 48,000 people in the U.S. died from suicide (CDC)
 - 10th leading cause of death overall
 - Has been top-12 every year since 1975
 - 2nd leading cause of death for ages 10-34
- Rates of suicide have been steadily increasing since 1999 (NIMH)
 - 745,356 U.S. deaths from 1999-2018
 - Issue across all demographics



Background on Suicide

- Suicide and suicide attempts are types of self-directed violence
 - Lifetime prevalence rate for self-directed violence ~6 percent
 - Estimates range from 1 in 9 to 1 in 25 suicide attempts result in suicide (Canner et al, 2018)
 - About 650,000 people are treated by hospital emergency departments each year after suicide attempts (SAMHSA)



Demographics of Suicide

- Highest rates of suicide occur among American Indians/Alaska Natives and White populations
- Ethnic minorities account for 15% of suicide deaths (Caetano et al., 2015)
- Suicide has increased in younger populations, however most deaths occur for those around middle-age/older adults
- Men experience a suicide death rate 4x higher than women, however women attempt suicide at a higher rate (AFSP, 2020)



Populations Most At-Risk (SAMHSA)

- Individuals who have attempted suicide previously
- Individuals who experience chronic pain
- American Indians/Alaska Natives
- Individuals who have lost someone to suicide
- Individuals in justice and child welfare settings
- Individuals who participate in non-suicidal self-injury
- Members of the armed forces as well as veterans
- Members of the LGBTQ+ community
- Men in midlife and older men
- Individuals who have a mental health and/or **substance use disorder(s)**



Linking Suicide to Substance Use

- Research has shown that substance use, in particular opioid or alcohol misuse, directly increases the risk of suicide (SAMHSA, 2016; CDC)
 - 22 percent of suicides involve alcohol intoxication
 - 20 percent of suicide involve opioids
- Both substance use and thoughts of suicide can stem from underlying mental health issues



Prevalence of Substance Use (SAMHSA)

In 2018, data from the United States found that:

- 1 in 12 individuals experienced a substance use disorder (SUD)
- 10.3 million people misused opioids
 - Most commonly prescription pain relievers, either on their own or in conjunction with heroin
- 5.4% of Americans experienced an alcohol use disorder
 - 67.1 million Americans were regular binge drinkers



Substance Use Fatalities (CDC)

- Between 1999 and 2018, the fatal drug-involved overdose rate increased 243%
- 16,849 fatal overdoses in 1999 compared to 67,367 in 2018
- Fatal opioid overdoses experienced the largest number of additional overdose deaths, from 8,050 in 1999 to 46,802 in 2018
- Alcohol-induced related deaths are estimated at anywhere from 35,800 to 88,000 every year



Substance Use and Suicide

- Suicide is the leading cause of death among individuals who are misusing substances
- Those with an SUD experience a 7 percent lifetime risk of a fatal suicide (Miller et al., 2016)
- Both suicide and substance use are connected to and impacted by adverse life experiences
- Substance use and suicide share many of the same risk factors



Alcohol and Suicide

- Alcohol has been closely associated with suicide, with acute alcohol intoxication is present 30%-40% of the time when an attempted suicide occurs (SAMSHA, 2016)
- Heavy alcohol drinkers have a 5-10 times higher risk of suicide
- Those who abuse alcohol experience a 10%-15% lifetime risk for suicide (Pompili, 2010)
- Impulse control, risky behavior, impaired cognitive abilities, and mental health issues can all be influenced by alcohol leading to a higher risk of suicidal behaviors



Opioids and Suicide

- Opioids cover both prescription medications as well as illicit substances, such as heroin
- Frequent opioid misuse (in the past and currently) can increase suicidal ideation, planning, and the number of attempts
- Older adults (age 50+) had an 84% increase in suicidal ideation if they were misusing prescription opioids (Schepis, 2019)
- Opioid overdose is the 2nd most common cause of death (16.2%) for individuals experiencing chronic pain (Ilgen, 2018)
 - One study found opioids present in 52% of chronic pain suicides



Opioids and Suicide

- Some research suggests that there may be an underreporting of opioid related suicides
 - Data from the U.S. National Violent Death Reporting System
 - Designating ‘suicide from drug intoxication’ as a person’s cause-of-death often requires meeting a significantly higher set of criteria than for other common causes of mortality
 - Most common form of attempted suicide is drug ingestion, highest fatalities with firearms and hanging



Summary of Evidence-Based Practices

- Because of the connections between substance use and suicide, there are often opportunities for collaboration between service providers
- But few EBPs *in prevention* exist to address both issues
 - Suicide prevention more commonly collaborates with substance use treatment
- However, the evidence that does exist suggests greater collaborations may be possible
 - Verified screening tools
 - Environmental strategies
 - Others?



Screening Tools

- Studies have identified screening tools effective at identifying people at-risk for substance use and at-risk for suicide
 - Addiction Brief RiskScale Assessment
 - Stratification Tool for Opioid Risk Mitigation
 - Reynolds Adolescent Depression Scale-2: Short Form & Alcohol Use Disorders Identification Test
 - Ask Suicide-Screening Questions & HEADS-ED
- Interactive Voice Recording Surveys can be effective methods of data collection



Collaborating with Screening Tools

- These screening tools can identify people at-risk for substance use and at-risk for suicide
 - Meaning they identify people who may benefit from substance use prevention and suicide prevention services
- Even if there is not a single EBP to address both needs, many EBPs to address either issue exist
- All service providers can benefit from expanded surveillance to identify people in need of services
- Prevention organizations could consider partnering with suicide prevention organizations to expand screening services



Environmental Strategies- Alcohol

- Research identified several environmental strategies significantly associated with reducing rates of alcohol misuse and of suicide, including:
 - Increasing alcohol taxation rates
 - Increasing the minimum legal drinking age
 - Decreasing density of alcohol retail outlets
 - Enforcing zero tolerance laws
 - Implementing alcohol misuse awareness campaigns
 - Municipalizing/nationalizing alcohol retail outlets
 - Regulating extremely high proof alcohol products



Environmental Strategies- Opioids

- Prescriber education significantly associated with reduced rates of opioid misuse, risk of suicide, and unintentional fatal overdose
- Prescription drug monitoring programs are significantly associated with reduced rates of opioids misuse, may reduce risk of suicide as well
- Many other strategies found to reduce rates of misuse and unintentional fatal overdose, but have not been assessed for potential suicide impact



Environmental Strategy Collaboration

- Environmental strategies are broad context changes that often require outside stakeholder support to enact
 - See February 25, 2020 PTTC webinar for more information on environmental strategies
- Substance use prevention and suicide prevention stakeholders could work together to educate and advocate for strategies mutually beneficial to their goals
 - Including developing materials (e.g. for opioid prescribers)



Other Opportunities for Collaboration

- Implementing EBPs is important; they give the greatest confidence that our efforts will be effective
- However, it is only through research, and pilot programs, that new EBPs can be identified and disseminated
- If your needs assessment finds that co-occurring risk of substance use and suicide is an issue in populations that you are serving, consider whether the capacity exists to test new initiatives



Evaluation

Your feedback is important!

Evaluation Form

To complete the evaluation form, go to <https://ttc-gpra.org/P?s=317583> or scan the QR code.



Evaluation Form
QR Code

Resource Page

Upon completion of the evaluation form, your browser will redirect to the resources page from today's webinar, which will include a PDF of the PowerPoint slides.

Certificate of Attendance / Contact Hour

For viewing the live webinar today, you can receive up to one (1) NADAAC contact hour of attendance. You may request a certificate from a link provided at the TOP of the RESOURCES PAGE. Certificates must be requested within 1 week of the webinar. Certificates will be processed within 30 days.

Recording

A recording of today's webinar will be available online at <http://bit.ly/ce-pttc> in 2-3 weeks.



Contact Us



Central East (HHS Region 3)

PTTC

Prevention Technology Transfer Center Network
Funded by Substance Abuse and Mental Health Services Administration

a program managed by



Central East PTTC website:
www.pttcnetwork.org/centraleast

Danya Institute website:
www.danyainstitute.org

Deborah Nixon-Hughes, Director:
dhughes@danyainstitute.org

Danya Institute email and phone:
info@danyainstitute.org
(240) 645-1145

Funding for this presentation was made possible by SAMHSA grant no. 1H79SP081018. The views expressed by speakers and moderators do not necessarily reflect the official policies of HHS; nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government.



Central East (HHS Region 3)

PTTC

